arXiv:1511.03308v2 [math.CA] 8 Dec 2015

GENERALIZED HERMITE-HADAMARD-FEJER TYPE INEQUALITIES FOR GA-CONVEX FUNCTIONS VIA FRACTIONAL INTEGRAL

İMDAT İŞCAN AND SERCAN TURHAN

Abstract. In this paper, it is a fuction that is a GA-convex differentiable for a new identity. As a result of this identity some new and general fractional integral inequalities for differentiable GA-convex functions are obtained.

1. Introduction

The classical or the usual convexity is defined as follows:
A function $f:\emptyset\neq I\subseteq\mathbb{R}\longrightarrow\mathbb{R}$, is said to be convex on I if inequality

$$f\left(tx+\left(1-t\right)y\right)\leq tf(x)+\left(1-t\right)f(y)$$

holds for all $x,y\in I$ and $t\in[0,1]$.

A number of papers have been written on inequalities using the classical convexty and one of the most fascinating inequalities in mathematical analysis is stated as follows

$$f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int_{a}^{b}f(x)dx\leq\frac{f(a)+f(b)}{2}, \tag{1.1}$$

where $f:I\subseteq\mathbb{R}\longrightarrow\mathbb{R}$ be a convex mapping and $a,b\in I$ with $a\leq b$. Both the inequalities hold in reversed direction if f is concave. The inequalities stated in (1.1) are known as Hermite-Hadamard inequalities.

For more results on (1.1) which provide new proof, significantly extensions, generalizations, refinements, counterparts, new Hermite-Hadamard-type inequalities and numerous applications, we refer the interested reader to [4]-[6],[8, 10, 11, 13, 14] and the references therein.

The usual notion of convex function have been generalized in diverse manners. One of them is the so called GA-convex functions and is stated in the defination below.

Definition 1. [10, 11] *A function $f:I\subseteq\mathbb{R}=(0,\infty)\longrightarrow\mathbb{R}$ is said to be GA-convex function on I if*

$$f\left(x^{\lambda}y^{1-\lambda}\right)\leq\lambda f\left(x\right)+\left(1-\lambda\right)f\left(y\right)$$

2000 *Mathematics Subject Classification.* Primary 26D15; Secondary 26A51, Third 26D10, Fourth 26A15.

Key words and phrases. GA-convex, Hermite-Hadamard-Fejer type inequalities, Fractional Integral.

holds for all $x, y \in I$ and $\lambda \in [0,1]$, where $x^{\lambda}y^{1-\lambda}$ and $\lambda f(x) + (1-\lambda) f(y)$ are respectively the weighted geometric mean of two positive numbers x and y and the weighted arithmetic mean of $f(x)$ and $f(y)$.

The defination of GA-convexity is further generalized as s-GA-convexity in the second sense as follows.

Definition 2. [17] *A function $f : I \subseteq \mathbb{R} = (0,\infty) \longrightarrow \mathbb{R}$ is said to be s-GA-convex function on I if*

$$f\left(x^{\lambda}y^{1-\lambda}\right) \leq \lambda^{s} f(x) + (1-\lambda)^{s} f(y)$$

holds for all $x, y \in I$ and $\lambda \in [0,1]$ and for some $s \in (0,1]$.

For the properties of GA-convex functions and GA-s-convex function, we refer the reader to [9, 12, 14, 15, 16, 17, 21, 22] and the reference there in.

Most recently, a number of findings have been seen on Hermite-Hadamard type integral inequalities for GA-convex and for GA-s-convex functions.

Zang at all. in [20] established the following Hermite-Hadamard type integral inequalities for GA-convex function.

Theorem 1. [20] *Let $f : I \subseteq \mathbb{R}_{+} \to \mathbb{R}$ be differentiable on I°, and $a, b \in I$ with $a < b$ and $f' \in L[a,b]$. If $|f'|^{q}$ is GA-convex on $[a,b]$ for $q \geq 1$, then*

$$\left| bf(b) - af(a) - \int_{a}^{b} f(x)dx \right| \leq \frac{\left[(b-a)A(a,b)\right]^{1-1/q}}{2^{1/q}} \tag{1.2}$$

$$\times \left\{ \left[L(a^2,b^2) - a^2\right] |f'(a)|^{q} + \left[b^2 - L(a^2,b^2)\right] |f'(b)|^{q} \right\}^{1/q}.$$

Theorem 2. [20] *Let $f : I \subseteq \mathbb{R}_{+} \to \mathbb{R}$ be differentiable on I°, and $a, b \in I$ with $a < b$ and $f' \in L[a,b]$. If $|f'|^{q}$ is GA-convex on $[a,b]$ for $q > 1$, then*

$$\left| bf(b) - af(a) - \int_{a}^{b} f(x)dx \right| \leq (\ln b - \ln a) \tag{1.3}$$

$$\times \left[L(a^{2q/(q-1)}, b^{2q/(q-1)}) - a^{2q/(q-1)} \right]^{1-1/q} \left[A\left(|f'(a)|^{q}, |f'(b)|^{q}\right) \right]^{1/q}.$$

Theorem 3. *Let $f : I \subseteq \mathbb{R}_{+} \to \mathbb{R}$ be differentiable on I°, and $a, b \in I$ with $a < b$ and $f' \in L[a,b]$. If $|f'|^{q}$ is GA-convex on $[a,b]$ for $q > 1$ and $2q > p > 0$, then*

$$\left| bf(b) - af(a) - \int_{a}^{b} f(x)dx \right| \leq \frac{(\ln b - \ln a)^{1-1/q}}{p^{1/q}} \tag{1.4}$$

$$\times \left[L(a^{(2q-p)/(q-1)}, b^{(2q-p)/(q-1)}) \right]^{1-1/q}$$

$$\times \left\{ \left[L(a^{p},b^{p}) - a^{p}\right] |f'(a)|^{q} + \left[b^{p} - L(a^{p},b^{p})\right] |f'(b)|^{q} \right\}^{1/q}.$$

Theorem 4. *Suppose that $f : I \subseteq \mathbb{R} = (0,\infty) \longrightarrow \mathbb{R}$ is s-GA-convex function in the second sense, where $s \in [0,1)$ and let $a, b \in [0,\infty)$, $a < b$. If $f \in L[a,b]$, then the following inequalities hold*

$$2^{s-1} f\left(\sqrt{ab}\right) \leq \frac{1}{lnb - lna} \int_{a}^{b} \frac{f(x)}{x} dx \leq \frac{f(a) + f(b)}{s+1}. \tag{1.5}$$

the constant $k = \frac{1}{s+1}$ *is the best possible in the second inequality in (1.1).*

If f in Theorem 5 is GA-convex function, then we get the following inequalities.

$$f\left(\sqrt{ab}\right) \leq \frac{1}{lnb - lna}\int_a^b \frac{f(x)}{x}dx \leq \frac{f(a)+f(b)}{2}. \tag{1.6}$$

For more results on GA-convex function and s-GA-convex function see e.g [9, 17, 21] and [22].

Definition 3. *A function* $f : I \subseteq \mathbb{R} = (0,\infty) \longrightarrow \mathbb{R}$ *is said to be geometrically symmetric with respect to* $\sqrt{ab}$ *if the inequality*

$$g\left(\frac{ab}{x}\right) = g\left(x\right)$$

holds for all $x \in [a,b]$

Definition 4. *Let* $f \in L[a,b]$. *The right-hand side and left-hand side Hadamard fractional integrals* $J^{\alpha}_{a^+}f$ *and* $J^{\alpha}_{b^-}f$ *of order* $\alpha > 0$ *with* $b > a \leq 0$ *are defined by*

$$J^{\alpha}_{a^+}f(x) = \frac{1}{\Gamma(\alpha)}\int_a^x \left(\ln\frac{x}{t}\right)^{\alpha-1} f(t)\frac{dt}{t},\ x > a$$

$$J^{\alpha}_{b^-}f(x) = \frac{1}{\Gamma(\alpha)}\int_x^b \left(\ln\frac{t}{x}\right)^{\alpha-1} f(t)\frac{dt}{t},\ x < b$$

respectively where $\Gamma(\alpha)$ *is the Gamma function defined by* $\Gamma(\alpha) = \int_0^{\infty} e^{-t}t^{\alpha-1}$

Lemma 1. *For* $0 < \theta \leq 1$ *and* $0 < a \leq b$ *we have*

$$\left|a^{\theta} - b^{\theta}\right| \leq (b-a)^{\theta}.$$

In [1] D. Y. Hwang found out a new identity and by using this identity, established a new inequalities. In this paper, we established a new identity similar to that's identity in [1] and then we obtained some new and general integral inequalities for differentiable GA-convex functions using this lemma.

2. Main result

Throughout this section, let $\|g\|_{\infty} = \sup_{x\in[a,b]}|g(x)|$, for the continuous function $g : [a,b] \longrightarrow \mathbb{R} = (0,\infty) \longrightarrow \mathbb{R}$ be differentiable mapping I^o, where $a, b \in I$ with $a \leq b$, and $h : [a,b] \longrightarrow [0,\infty)$ be differentiable mapping. In the following sections, for convenience, let the notion $L(t) = a^t G^{1-t}$, $U(t) = b^t G^{1-t}$ and $G = G(a,b) = \sqrt{ab}$.

Lemma 2. *If* $f' \in L(a,b)$ *then the following inequality holds:*

$$[h(b) - 2h(a)]\frac{f(a)}{2} + h(b)\frac{f(b)}{2} - \int_a^b f(x)h'(x)dx \tag{2.1}$$

$$= \frac{lnb - lna}{4} \left\{ \int_0^1 \left[2h\left(a^t G^{1-t}\right) - h(b)\right] f'\left(a^t G^{1-t}\right) a^t G^{1-t} dt \right.$$
$$\left. + \int_0^1 \left[2h\left(b^t G^{1-t}\right) - h(b)\right] f'\left(b^t G^{1-t}\right) b^t G^{1-t} dt \right\}.$$

Proof. By the integration by parts, we have

$$I_1 = \int_0^1 \left[2h\left(a^t G^{1-t}\right) - h(b)\right] d\left(f\left(a^t G^{1-t}\right)\right)$$
$$= \left[2h\left(a^t G^{1-t}\right) - h(b)\right] f\left(a^t G^{1-t}\right)\Big|_0^1$$
$$-2\ln\left(\frac{a}{G}\right) \int_0^1 f\left(a^t G^{1-t}\right) h'\left(a^t G^{1-t}\right) a^t G^{1-t} dt$$

and

$$I_2 = \int_0^1 \left[2h\left(b^t G^{1-t}\right) - h(b)\right] d\left(f\left(b^t G^{1-t}\right)\right)$$
$$= \left[2h\left(b^t G^{1-t}\right) - h(b)\right] f\left(b^t G^{1-t}\right)\Big|_0^1$$
$$-2\ln\left(\frac{b}{G}\right) \int_0^1 f\left(a^t G^{1-t}\right) h'\left(b^t G^{1-t}\right) b^t G^{1-t} dt$$

Therefore

(2.2)
$$\frac{I_1 + I_2}{2} = [h(b) - 2h(a)] \frac{f(a)}{2} + h(b)\frac{f(b)}{2} - \frac{lnb - lna}{2} \left\{ \int_0^1 f\left(a^t G^{1-t}\right) h'\left(a^t G^{1-t}\right) a^t G^{1-t} dt \right.$$
$$\left. + \int_0^1 f\left(a^t G^{1-t}\right) h'\left(a^t G^{1-t}\right) a^t G^{1-t} dt \right\}$$

This complete the proof □

Lemma 3. *For $a, G, b > 0$, we have*

$$\zeta_1(a,b) = \int_0^1 t a^t G^{1-t} \left|2h\left(a^t G^{1-t}\right) - h(b)\right| dt \tag{2.3}$$

(2.4)
$$\zeta_2(a,b) = \int_0^1 (1-t) a^t G^{1-t} \left|2h\left(a^t G^{1-t}\right) - h(b)\right| dt + \int_0^1 (1-t) b^t G^{1-t} \left|2h\left(b^t G^{1-t}\right) - h(b)\right| dt$$

$$\zeta_3(a,b)=\int_0^1 tb^tG^{1-t}\left|2h\left(b^tG^{1-t}\right)-h(b)\right|dt \tag{2.5}$$

Theorem 5. *Let $f:I\subseteq\mathbb{R}=(0,\infty)\longrightarrow\mathbb{R}$ be differentiable mapping I^o, where $a,b\in I^o$ with $a<b$. If the mapping $|f'|$ is GA-convex on $[a,b]$, then the following inequality holds:*

$$\left|[h(b)-2h(a)]\frac{f(a)}{2}+h(b)\frac{f(b)}{2}-\int_a^b f(x)h'(x)dx\right| \tag{2.6}$$
$$\leq\frac{\ln b-\ln a}{4}\left[\zeta_1(a,b)\left|f'(a)\right|+\zeta_2(a,b)\left|f'(G)\right|+\zeta_3(a,b)\left|f'(b)\right|\right]$$

where $\zeta_1(a,b),\zeta_2(a,b),\zeta_3(a,b)$ are defined in Lemma 2.

Proof. Continuing inequality (2.1) in Lemma 1

$$\left|[h(b)-2h(a)]\frac{f(a)}{2}+h(b)\frac{f(b)}{2}-\int_a^b f(x)h'(x)dx\right| \tag{2.7}$$
$$\leq\frac{lnb-lna}{4}\left\{\int_0^1\left|2h\left(a^tG^{1-t}\right)-h(b)\right|\left|f'\left(a^tG^{1-t}\right)a^tG^{1-t}\right|dt\right.$$
$$\left.+\int_0^1\left|2h\left(b^tG^{1-t}\right)-h(b)\right|\left|f'\left(b^tG^{1-t}\right)b^tG^{1-t}\right|dt\right\}$$

Using $|f'|$ is GA-convex in (2.7)

$$\left|[h(b)-2h(a)]\frac{f(a)}{2}+h(b)\frac{f(b)}{2}-\int_a^b f(x)h'(x)dx\right| \tag{2.8}$$
$$\leq\frac{lnb-lna}{4}\left\{\int_0^1\left|2h\left(a^tG^{1-t}\right)-h(b)\right|\left[t\left|f'(a)\right|+(1-t)\left|f'(G)\right|\right]a^tG^{1-t}dt\right.$$
$$\left.+\int_0^1\left|2h\left(b^tG^{1-t}\right)-h(b)\right|\left[t\left|f'(b)\right|+(1-t)\left|f'(G)\right|\right]dt\right\},$$

by (2.8) and Lemma 2, this proof is complete. □

Corollary 1. *$g:[a,b]\longrightarrow[0,\infty)$ be continuous positive mapping and symmetric to $\sqrt{ab}$. Let $h(x)=\int_a^x\left[\left(ln\frac{b}{t}\right)^{\alpha-1}+\left(ln\frac{t}{a}\right)^{\alpha-1}\right]\frac{g(t)}{t}dt$ for all $t\in[a,b]$ and $\alpha>0$ in Teorem 5, we obtain:*

$$\left|\left(\frac{f(a)+f(b)}{2}\right)\left[J^\alpha_{a^+}g(b)+J^\alpha_{b^-}g(a)\right]-\left[J^\alpha_{a^+}(fg)(b)+J^\alpha_{b^-}(fg)(a)\right]\right| \tag{2.9}$$

$$\leq \frac{(\ln b-\ln a)^{\alpha+1}}{2^{\alpha+1}\Gamma(\alpha+1)}\|g\|_{\infty}\left[C_1(\alpha)|f'(a)|+C_2(\alpha)|f'(G)|+C_3(\alpha)|f'(b)|\right]$$

where

$$C_1(\alpha)=\int_0^1\left[(1+t)^{\alpha}-(1-t)^{\alpha}\right]ta^tG^{1-t}dt$$

$$C_2(\alpha)=\int_0^1(1-t)\left[(1+t)^{\alpha}-(1-t)^{\alpha}\right]\left[a^tG^{1-t}+b^tG^{1-t}\right]dt$$

$$C_3(\alpha)=\int_0^1\left[(1+t)^{\alpha}-(1-t)^{\alpha}\right]tb^tG^{1-t}dt$$

Specially in (2.9) and using Lemma 2, for $0<\alpha\leq 1$ *we have:*

$$\left|\left(\frac{f(a)+f(b)}{2}\right)\left[J^{\alpha}_{a^+}g(b)+J^{\alpha}_{b^-}g(a)\right]-\left[J^{\alpha}_{a^+}(fg)(b)+J^{\alpha}_{b^-}(fg)(a)\right]\right| \tag{2.10}$$

$$\leq \frac{(\ln b-\ln a)^{\alpha+1}}{2\Gamma(\alpha+1)}\|g\|_{\infty}\left[C_1(\alpha)|f'(a)|+C_2(\alpha)|f'(G)|+C_3(\alpha)|f'(b)|\right]$$

where

$$C_1(\alpha)=\int_0^1 t^{\alpha+1}a^tG^{1-t}$$

$$C_2(\alpha)=\int_0^1\left[(1-t)\,t^{\alpha}a^tG^{1-t}+(1-t)\,t^{\alpha}b^tG^{1-t}\right]dt$$

$$C_3(\alpha)=\int_0^1 t^{\alpha+1}b^tG^{1-t}$$

Proof. By left side of inequality (2.8) in Teorem 5, when we write $h(x)=\int_a^x\left[\left(ln\frac{b}{t}\right)^{\alpha-1}+\left(ln\frac{t}{a}\right)^{\alpha-1}\right]\frac{g(t)}{t}dt$ for all $t\in[a,b]$, we have

$$\left|\Gamma(\alpha)\left(\frac{f(a)+f(b)}{2}\right)\left[J^{\alpha}_{a^+}g(b)+J^{\alpha}_{b^-}g(a)\right]-\Gamma(\alpha)\left[J^{\alpha}_{a^+}(fg)(b)+J^{\alpha}_{b^-}(fg)(a)\right]\right|$$

.

On the other hand, right side of inequality (2.8)

(2.11)

$$\leq \frac{lnb-lna}{4}\left\{\int_0^1\left|\begin{array}{c}2\int_a^{a^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\\ -\int_a^b\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\end{array}\right|\left[t|f'(a)|+(1-t)|f'(G)|\right]a^tG^{1-t}dt\right.$$

$$+\int_0^1 \left| \begin{array}{l} 2\int_a^{b^tG^{1-t}} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx \\ -\int_a^{b} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx \end{array} \right| [t\,|f'(b)|+(1-t)\,|f'(G)|]\,b^tG^{1-t}dt \Bigg\}$$

.

Since $g(x)$ is symmetric to $x=\sqrt{ab}$, we have

(2.12)
$$\left|2\int_a^{a^tG^{1-t}} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx-\int_a^{b} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|$$
$$=\left|\int_{a^tG^{1-t}}^{b^tG^{1-t}} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|$$

and

(2.13)
$$\left|2\int_a^{b^tG^{1-t}} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx-\int_a^{b} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|$$
$$=\left|\int_{a^tG^{1-t}}^{b^tG^{1-t}} \left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|$$

for all $t\in[0,1]$.

By (2.10), (2.11) and (2.12), we have

$$\left|\left(\frac{f(a)+f(b)}{2}\right)\left[J^{\alpha}_{a^+}g(b)+J^{\alpha}_{b^-}g(a)\right]-\left[J^{\alpha}_{a^+}\left(fg\right)(b)+J^{\alpha}_{b^-}\left(fg\right)(a)\right]\right| \tag{2.14}$$
$$\leq \frac{lnb-lna}{4\Gamma(\alpha)}\left\{\int_0^1\left|\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|[t\,|f'(a)|+(1-t)\,|f'(G)|]\,a^tG^{1-t}dt\right.$$
$$\left.+\int_0^1\left|\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{g(x)}{x}dx\right|[t\,|f'(b)|+(1-t)\,|f'(G)|]\,b^tG^{1-t}dt\right\}$$
$$\leq \frac{lnb-lna}{4\Gamma(\alpha)}\|g\|_{\infty}\left\{\int_0^1\left[\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{1}{x}dx\right][t\,|f'(a)|+(1-t)\,|f'(G)|]\,a^tG^{1-t}dt\right.$$
$$\left.+\int_0^1\left[\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{1}{x}dx\right][t\,|f'(b)|+(1-t)\,|f'(G)|]\,b^tG^{1-t}dt\right\}.$$

In the last inequality,

(2.15)
$$\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{1}{x}dx=\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left(\ln\frac{b}{x}\right)^{\alpha-1}\frac{1}{x}dx+\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left(\ln\frac{x}{a}\right)^{\alpha-1}\frac{1}{x}dx$$
$$=\frac{2.(\ln b-\ln a)^{\alpha}}{2^{\alpha}.\alpha}\left[(1+t)^{\alpha}-(1-t)^{\alpha}\right].$$

By Lemma 1, we have

$$\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left[\left(\ln\frac{b}{x}\right)^{\alpha-1}+\left(\ln\frac{x}{a}\right)^{\alpha-1}\right]\frac{1}{x}dx=\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left(\ln\frac{b}{x}\right)^{\alpha-1}\frac{1}{x}dx+\int_{a^tG^{1-t}}^{b^tG^{1-t}}\left(\ln\frac{x}{a}\right)^{\alpha-1}\frac{1}{x}dx$$
$$\leq\frac{2.(\ln b-\ln a)^{\alpha}}{\alpha}t^{\alpha}$$

A combination of (2.13) and (2.14), we have (2.9). This complete is proof. □

Corollary 2. *In Corollary 1,*

(1)If $\alpha=1$ is in corollary, we obtain following Hermite-Hadamard-Fejer Type inequality for GA-convex function which is related in (2.10):

$$\left|\left[\frac{f(a)+f(b)}{2}\right]\int_a^b\frac{g(x)}{x}dx-\int_a^b f(x)\frac{g(x)}{x}dx\right|\leq \tag{2.16}$$
$$\frac{(\ln b-\ln a)^2}{4}\|g\|_{\infty}\left[C_1(1)\left|f'(a)\right|+C_2(1)\left|f'(G)\right|+C_3(1)\left|f'(b)\right|\right]$$

where for $a,b,G>0$, we have

$$C_1(1)=\int_0^1 t^2a^tG^{1-t}dt=\frac{2}{\ln b-\ln a}\left\{-a-\frac{4a}{\ln b-\ln a}-\frac{8a-8G}{(\ln b-\ln a)^2}\right\},$$

$$C_2(1)=\int_0^1 t(1-t)a^tG^{1-t}dt+\int_0^1 t(1-t)b^tG^{1-t}dt=\frac{2}{\ln b-\ln a}\left\{\frac{2(a+b+2G)}{\ln b-\ln a}+\frac{8(a-b)}{(\ln b-\ln a)^2}\right\},$$

and

$$C_3(1)=\int_0^1 t^2a^tG^{1-t}dt=\frac{2}{\ln b-\ln a}\left\{b-\frac{4b}{\ln b-\ln a}+\frac{8b-8G}{(\ln b-\ln a)^2}\right\}.$$

(2)If $g(x)=1$ is in corollary, we obtain following Hermite-Hadamard-Fejer Type inequality for GA-convex function which is related in (2.9):

$$\left|\left(\frac{f(a)+f(b)}{2}\right)-\frac{\Gamma(\alpha+1)}{2(\ln b-\ln a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right| \tag{2.17}$$
$$\leq\frac{(\ln b-\ln a)}{2^{\alpha+2}}\left[C_1(\alpha)\left|f'(a)\right|+C_2(\alpha)\left|f'(G)\right|+C_3(\alpha)\left|f'(b)\right|\right].$$

(3)If $g(x)=1$ and $\alpha=1$ is in corollary, we obtain following Hermite-Hadamard-Fejer Type inequality for GA-convex function which is related in (2.10):

$$\left|\left(\frac{f(a)+f(b)}{2}\right)-\frac{1}{(\ln b-\ln a)}\int_a^b\frac{f(x)}{x}dx\right| \tag{2.18}$$

$$\leq\frac{(\ln b-\ln a)}{4}\left[C_1(1)\left|f'(a)\right|+C_2(1)\left|f'(G)\right|+C_3(1)\left|f'(b)\right|\right].$$

Theorem 6. *Let $f:I\subseteq\mathbb{R}^+=(0,\infty)\longrightarrow\mathbb{R}$ be differentiable mapping I^o, where $a,b\in I$ with $a<b$, and $g:[a,b]\longrightarrow[0,\infty)$ be continuous positive mapping and symmetric to $\sqrt{ab}$ and $\frac{1}{q}+\frac{1}{p}=1$. If the mapping $|f'|^q$ is GA-convex on $[a,b]$, then the following inequality holds:*

$$\left|[h(b)-2h(a)]\frac{f(a)}{2}+h(b)\frac{f(b)}{2}-\int_a^b f(x)h'(x)dx\right| \tag{2.19}$$

$$\leq\frac{\ln b-\ln a}{4}\left\{\begin{array}{c}\left(\int_0^1\left|2h(a^tG^{1-t})-h(b)\right|dt\right)^{1-\frac{1}{q}}\times\\ \left(\int_0^1\begin{array}{c}\left(\left|2h(a^tG^{1-t})-h(b)\right|dt\right)\\ \times\left(ta^{qt}G^{q(1-t)}\left|f'(a)\right|^q+(1-t)a^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right)\end{array}\right)^{\frac{1}{q}}\\ +\left(\int_0^1\left|2h(b^tG^{1-t})-h(b)\right|dt\right)^{1-\frac{1}{q}}\times\\ \left(\int_0^1\begin{array}{c}\left(\left|2h(b^tG^{1-t})-h(b)\right|dt\right)\\ \times\left(tb^{qt}G^{q(1-t)}\left|f'(b)\right|^q+(1-t)b^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right)\end{array}\right)^{\frac{1}{q}}\end{array}\right\}$$

Proof. Continuing from (2.7) in Theorem 5, we use Holder Inequality and we use that $|f'|^q$ is GA-convex. Thus this proof is complete. □

Corollary 3. *Let $h(x)=\int_a^x\left[\left(ln\frac{b}{t}\right)^{\alpha-1}+\left(ln\frac{t}{a}\right)^{\alpha-1}\right]\frac{g(t)}{t}dt$ for all $t\in[a,b]$ in Teorem 6, we obtain:*

$$\left|\left(\frac{f(a)+f(b)}{2}\right)\left[J^\alpha_{a^+}g(b)+J^\alpha_{b^-}g(a)\right]-\left[J^\alpha_{a^+}(fg)(b)+J^\alpha_{b^-}(fg)(a)\right]\right| \tag{2.20}$$

$$\leq\frac{(\ln b-\ln a)^{\alpha+1}\left\|g\right\|_\infty}{2^{\alpha+1}\Gamma(\alpha+1)}(\frac{2^{\alpha+2}-2^2}{\alpha+1})^{1-\frac{1}{q}}\left[C_1(\alpha,q)\left|f'(a)\right|^q+C_2(\alpha,q)\left|f'(G)\right|^q+C_3(\alpha,q)\left|f'(b)\right|^q\right]^{\frac{1}{q}}$$

where for $q>1$

$$C_1(\alpha,q)=\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]ta^{qt}G^{q(1-t)}dt$$

$$C_2(\alpha,q)=\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right](1-t)\left(a^{qt}G^{q(1-t)}+b^{qt}G^{q(1-t)}\right)dt$$

$$C_3(\alpha,q)=\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]tb^{qt}G^{q(1-t)}dt.$$

Proof. Continuing from (2.15) of Corollary 1 and (2.19) in Theorem 6,

$$\left|\left(\frac{f(a)+f(b)}{2}\right)\left[J^\alpha_{a^+}g(b)+J^\alpha_{b^-}g(a)\right]-\left[J^\alpha_{a^+}(fg)(b)+J^\alpha_{b^-}(fg)(a)\right]\right| \tag{2.21}$$

$$\leq\frac{(\ln b-\ln a)^{\alpha+1}\|g\|_\infty}{2^{\alpha+1}\Gamma(\alpha+1)}\left\{\begin{array}{c}\left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]dt\right)^{1-\frac{1}{q}}\times\\ \left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]\left(ta^{qt}G^{q(1-t)}\left|f'(a)\right|^q+(1-t)a^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right)dt\right)^{\frac{1}{q}}\\ +\\ \left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]dt\right)^{1-\frac{1}{q}}\times\\ \left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]\left(tb^{qt}G^{q(1-t)}\left|f'(b)\right|^q+(1-t)b^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right)dt\right)^{\frac{1}{q}}\end{array}\right\}$$

$$\leq\frac{(\ln b-\ln a)^{\alpha+1}\|g\|_\infty}{2^{\alpha+1}\Gamma(\alpha+1)}\left(\frac{2^{\alpha+1}-2}{\alpha+1}\right)^{1-\frac{1}{q}}\left[\begin{array}{c}\left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]\times\right.\\ \left.\left[ta^{qt}G^{q(1-t)}\left|f'(a)\right|^q+(1-t)a^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right]\ dt\right)^{\frac{1}{q}}\\ +\left(\int_0^1\left[(1+t)^\alpha-(1-t)^\alpha\right]\times\right.\\ \left.\left[tb^{qt}G^{q(1-t)}\left|f'(b)\right|^q+(1-t)b^{qt}G^{q(1-t)}\left|f'(G)\right|^q\right]\ dt\right)^{\frac{1}{q}}\end{array}\right]$$

By the power-mean inequality $\left(a^r+b^r<2^{1-r}(a+b)^r\ for\quad a>0,b>0,\quad r<1\right)$ and $\frac{1}{q}+\frac{1}{q}=1$ we have

$$\leq\frac{(\ln b-\ln a)^{\alpha+1}\|g\|_\infty}{2^{\alpha+1}\Gamma(\alpha+1)}\left(\frac{2^{\alpha+2}-2^2}{\alpha+1}\right)^{1-\frac{1}{q}} \tag{2.22}$$

$$\times\left[\int_0^1\left(\begin{array}{c}\left[(1+t)^\alpha-(1-t)^\alpha\right]ta^{qt}G^{q(1-t)}\left|f'(a)\right|^q+\\ \left[(1+t)^\alpha-(1-t)^\alpha\right](1-t)\left(\begin{array}{c}a^{qt}G^{q(1-t)}\\ +b^{qt}G^{q(1-t)}\end{array}\right)\left|f'(G)\right|^q\\ +\left[(1+t)^\alpha-(1-t)^\alpha\right]tb^{qt}G^{q(1-t)}\left|f'(b)\right|^q\end{array}\right)dt\right]^{\frac{1}{q}}$$

□

Corollary 4. *When $\alpha=1$ and $g(x)=\frac{1}{\ln b-\ln a}$ is taken in Corollary 3, we obtain:*

$$\left|\left(\frac{f(a)+f(b)}{2}\right)-\frac{1}{\ln b-\ln a}\int_a^b\frac{f(x)}{x}dx\right| \tag{2.23}$$

$$\leq\frac{(\ln b-\ln a)}{2^{1+\frac{1}{q}}}\left[C_1(1,q)\left|f'(a)\right|^q+C_2(1,q)\left|f'(G)\right|^q+C_3(1,q)\left|f'(b)\right|^q\right]^{\frac{1}{q}}.$$

This proof is complete.

References

[1] D-Y. Hwang, Some inequalities for differentiable convex mapping with application to weighted trapezoidal formula and higher moments of random variables, Applied Mathematics and Computation, 217 (2011), 9598-9605.

[2] D-Y. Hwang, Some inequalities for differentiable convex mapping with application to weighted midpoint formula and higher moments of random variables, Applied Mathematics and Computation, 232 (2014), 68-75.

[3] P.S. Bullen, Handbook of Means and Their Inequalities, Mathematics and its Applications, Volume 560, Kluwer Academic Publishers,Dordrecht/Boston/London, 2003.

[4] S.S. Dragomir, Hermite-Hadamard's type inequalities for convex funtions of selfadjoint operators in Hilbert spaces, Linear Algebra Appl. 436 (2012), no. 5, 1503-1515.

[5] S.S. Dragomir and C.E.M. Pearce, Selected topics on Hermite-Hadamard type inequalities and applications, RGMIA Monographs, 2000. Available online at http://rgmia.vu.edu.au/monographs/hermite hadamard.html.

[6] J. Hua, B.-Y. Xi, and F. Qi, Hermite-Hadamard type inequalities for geometrically s-convex functions, Commun.Korean Math.Soc.29 (2014), No.1, pp.51-63.

[7] H.Hudzik and Maligranda, Some remarks on s-convex functions, Aequationes Math. 48 (1994), no. 1,100-111. V. N. Huy and N. T. Chung, Some generalization of the Fejer and Hermite-Hadamard inequalities in Hölder space, J. Appl. Math. Inform. 29 (2011), no. 3-4, 859-868.

[8] I. Iscan, New estimates on generalization of some integral inequalities for s-convex functions and their applications, International Journal of Pure and Applied Mathematics, 86**,** No.4 (2013).

[9] I. Iscan, Hermite-Hadamard type inequalities for GA-s-convex functions, Le Matematiche,LXIX (2014)-Fasc. II, pp. 129-146.

[10] C. P. Niculescu, Convexity according to the geometric mean, Math. Inequal. Appl. 3 (2) (2000), 155–167. Available online at http://dx.doi.org/10.7153/mia-03-19.

[11] C. P. Niculescu, Convexity according to means, Math. Inequal. Appl. 6 (4) (2003), 571–579. Available online at http://dx.doi.org/10.7153/mia-06-53.

[12] M.A. Latif, New Hermite Hadamard type integral inequalities for GA-convex functions with applications. Volume 34, Issue 4 (Nov 2014).

[13] Y. Shuang, H.-P. Yin, and F. Qi, Hermite-Hadamard type integral inequalities for geometric-arithmetically s-convex functions, Analysis (Munich) 33 (2) (2013), 197-208. Available online at http://dx.doi.org/10.1524/anly.2013.1192.

[14] T.-Y. Zhang, A.-P. Ji and F. Qi, Some inequalities of Hermite-Hadamard type for GA-convex functions with applications to means, Le Matematiche, Vol. LXVIII (2013) – Fasc. I, pp. 229–239. doi: 10.4418/2013.68.1.17

[15] X.-M. Zhang, Y.-M. Chu, and X.-H. Zhang, The Hermite-Hadamard Type Inequality of GA-Convex Functions and Its Application, Journal of Inequalities and Applications, Volume 2010, Article ID 507560, 11 pages. doi:10.1155/2010/507560.

[16] A. P. Ji, T. Y. Zhang, F. Qi, Integral Inequalities of Hermite Hadamard Type (α, m)-GA convex Functions, Journal of Function Space and Applications, 2013 (2013), Article ID 823856, 8 pages.

[17] I. Iscan, Hermite Hadamard type inequalities for GA-s-convex functions, LE Mathematiche, LXIX (2014)-Fasc. II, pp. 129-146.

[18] C. P. Niculescu, Convexity according to geometric mean, Math. Inequal. Appl. 3 (2000), no. 2, 155-167.

[19] C. P. Niculescu, Convexity according to means, Math. Inequal. Appl. 3 (2003), no. 4, 571-579.

[20] Some inequalities of Hermite-Hadamard type for GA-Convex functions with applications to means. Le Matematiche, 48 (2013), no. 1, 229-239.

[21] M.A. Latif, S. S. Dragomir, E. Momoniat, Fejér type integral inequalities related with geometrically-arithmetically-convex functions with applications, (Submitted).

[22] M.A. Latif, S. S. Dragomir, E. Momoniat, Some Fejér type integral inequalities related with geometrically-arithmetically-convex functions with applications, (Submitted).

Department of Mathematics, Faculty of Arts and Sciences,, Giresun University, 28100, Giresun, Turkey.
E-mail address: imdati@yahoo.com, imdat.iscan@giresun.edu.tr

Dereli Vocational High School,, Giresun University, 28100, Giresun, Turkey.
E-mail address: sercanturhan28@gmail.com, imdat.sercan.turhan@giresun.edu.tr